

FarmLens Ltd

Website: farmlens.africa | App: app.farmlens.africa | Headquarters: Nairobi, Kenya



Crop details

Green Pea

Pisum sativum

Family: Fabaceae

Categories

Legumes & Pulses

Vegetables

Generated: 2026-06-03 13:01

Quick stats

Family	Fabaceae
Typical harvest	6.0 t/ha
Varieties	1
Pests and diseases	2
Seasons	1

Crop profile

Growth habit	annual
Days to harvest	80
Main uses	Fresh pods and shelled peas for food and processing.
Pollination	insect
Origin and where it grows	Common in cool East African highland vegetable production.

Weather, soil and spacing

Best temperature	10 - 22 °C
Rainfall	450 - 800 mm/yr
Altitude	800 - 2800 m
Best pH	6 - 7
Soil type	Fertile well-drained loam rich in organic matter.
Row spacing	40 cm
Plant spacing	20 cm
Planting depth	1.5 cm
Seed rate	4 kg/ha

Simple notes for farmers

About the crop: This crop is annual; it grows and is harvested in one season. Harvest typically starts about 80 days after planting.

Main use: Farmers mostly grow this crop for fresh pods and shelled peas for food and processing..

Pollination: Mainly insect; healthy flowers and pollinators improve fruit set.

Where it grows: Common in cool East African highland vegetable production.. Grouped under: Legumes & Pulses, Vegetables.

Best climate: 10 - 22 °C; 450 - 800 mm/yr; up to about 2800 m a.s.l.

Soil: Best at pH 6 - 7; fertile well-drained loam rich in organic matter..

Farmer guide (Mwongozo wa Mkulima)

<u>Planting</u>	Establish Green Pea in a fine weed-free seedbed and keep emergence moisture steady.
<u>Transplanting</u>	Direct seed or transplant depending on production system.
<u>Irrigation</u>	Maintain even soil moisture for steady Green Pea growth and quality.
<u>Fertigation</u>	Use split nitrogen and potassium for market-quality Green Pea.
<u>Pest scouting</u>	Scout Green Pea weekly for chewing pests, sap suckers, and foliar diseases.
<u>Pruning and training</u>	No pruning required unless sanitation or staking is needed.
<u>Harvest</u>	Harvest Green Pea at market maturity for the intended use.
<u>Postharvest</u>	Cool and shade Green Pea promptly after harvest.

Nutrient schedule (Mbolea kwa Hatua)

<u>#</u>	<u>Stage</u>	<u>DAP</u>	<u>Product</u>	<u>Rate</u>	<u>Targets (kg/ha)</u>	<u>Notes</u>
1	Basal	0	NPK 17-17-17	200 kg/ha	N: 34, P ₂ O ₅ : 34, K ₂ O: 34	Basal fertilizer for Green Pea.
2	Topdress	21	CAN	100 kg/ha	N: 26, P ₂ O ₅ : N/A, K ₂ O: N/A	Support active Green Pea vegetative growth.

Nutrient requirements

<u>Nutrient</u>	<u>Stage</u>	<u>Amount</u>	<u>Unit</u>
N	Basal	35	kg/ha
P ₂ O ₅	Basal	30	kg/ha
K ₂ O	Basal	35	kg/ha
N	Topdress	25	kg/ha
K ₂ O	Topdress	20	kg/ha

Field images



Varieties

<u>Name</u>	<u>Country</u>	<u>Maturity (days)</u>	<u>Traits</u>
Sugar Ann	KE	80	Early tender pods for fresh markets.

Fertilizer recommendations

<u>Stage</u>	<u>Product</u>	<u>Rate</u>	<u>Notes</u>
Planting	Well-rotted manure	5000	Improve soil structure before Green Pea planting.
Vegetative growth	CAN	100	Split topdress for Green Pea production.

Pests and diseases

<u>Name</u>	<u>Type</u>	<u>Symptoms</u>	<u>Management</u>
Aphids	pest	Leaf curling and sticky honeydew.	Scout frequently and use selective control when pressure builds.
Leaf spot complex	disease	Necrotic spotting and reduced leaf quality.	Improve airflow, rotate crops, and avoid prolonged leaf wetness.

Yields

<u>System</u>	<u>Typical</u>	<u>Min</u>	<u>Max</u>	<u>Notes</u>
Managed fresh-market production	6	4.2	9	Typical marketable Green Pea yield under irrigated or well-managed conditions.

Season calendars

<u>Country</u>	<u>Region</u>	<u>Planting</u>	<u>Harvest</u>
KE	Highland Vegetable Zones	Mar-Apr or Oct-Nov	Year-round depending on irrigation

Region suitability

<u>Country</u>	<u>Region</u>	<u>Suitability</u>
KE	Highland Vegetable Zones	High

Source: FarmLens Ltd - farmlens.africa and app.farmlens.africa. Headquarters: Nairobi, Kenya. This guide was generated from the FarmLens database.